

# MoorMan's® IGR Mineral® Block

To prevent the breeding of horn flies in the manure of treated cattle only. A pesticidally active mineral feed for beef cattle containing methoprene\* insect growth regulator for continuous feeding during the fly season.

## ACTIVE INGREDIENT:

S-methoprene (CAS Number 65733-16-6)..... 0.02%

OTHER INGREDIENTS: ..... 99.98%

Total 100.00%

## GUARANTEED ANALYSIS

Calcium (Ca), Min. ....13.2%, Max. .... 15.8%  
Phosphorus (P), Min. .... 7.0%  
Salt (NaCl), Min. ....25.0%, Max. .... 30.0%  
Iodine (I), Min. .... 0.015%  
Magnesium (Mg), Min. .... 0.2%  
Potassium (K), Min. .... 0.2%  
Copper (Cu), Min. .... 1100 ppm  
Selenium (Se), Min. .... 39 ppm  
Zinc (Zn), Min. .... 3800 ppm

Vitamin A, not less than 200,000 IU per pound.

## FEED INGREDIENTS

Defluorinated Phosphate, Salt, Soybean Meal,  
Monocalcium Phosphate, Dicalcium Phosphate,  
Calcium Carbonate, Cane Molasses, Iron Oxide,  
Manganous Oxide, Zinc Oxide, Lignin Sulfonate,  
Copper Sulfate, Vitamin E Supplement, Ferrous  
Sulfate, Magnesium Oxide, Vitamin A Acetate,  
Ethylenediamine Dihydriodide, Potassium Iodide,  
Sodium Selenite, Cobalt Carbonate, Mineral Oil,  
Cholecalciferol (Source of Vitamin D<sub>3</sub>).

Use this product as specified on the label.

Keep out of reach of children.

## CAUTION

EPA REG. NO. 1157-42

EPA Est. 1157-NE-002

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Make sure cattle are not starved for minerals or salt by providing them before feeding MoorMan's IGR Mineral Block. Then remove salt or mineral supplement and place MoorMan's IGR Mineral Block before cattle as the only source of minerals. Start feeding MoorMan's IGR Mineral Block continuously in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity.

The recommended consumption of MoorMan's IGR Mineral Block is 0.25 lb (1/4 lb) to 0.5 lb (1/2 lb) per 100 lb of body weight per month. Allow one MoorMan's IGR Mineral Block for each 15 head and place near watering or loafing areas. Protect blocks from rain.

If consumption of MoorMan's IGR Mineral Block is above 0.5 lb (1/2 lb) per 100 lb of body weight per month, reduce the number of feeding locations where this block is fed or relocate blocks in areas less frequented by cattle.

If consumption of MoorMan's IGR Mineral Block is below 0.25 lb (1/4 lb) per 100 lb of body weight per month, increase the number of feeding locations where this block is fed or relocate blocks in areas more frequented by cattle.

## STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry area. Do not stack anything on top of this block without placing a slip sheet or wooden pallet between blocks.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Completely empty container. Dispose of container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ADM Animal Nutrition, a division of Archer Daniels Midland Company, Quincy, IL 62305-3115

PRODUCT NO. 650AU

NET WT 50 LB (22.67 kg)

18517-CL

BLOCK

650AU

LOT NUMBER:



7 88399 60002 4