



Farmers Cooperative Association, Inc.

Medicated 16% Goat Textured

MEDICATED

For the prevention of coccidiosis caused by *Eimeria Crandallis*, *Eimeria Christenseni* and *Eimeria Ninahohlyakimovae*

ACTIVE DRUG INGREDIENT

Monensin,min. 20 g/ton

GUARANTEED ANALYSIS

Crude Protein,min. 16.00%
 (This product includes not more 1.3% equivalent crude protein from non-protein nitrogen)

Crude Fat,min. 2.25%
 Crude Fiber,max. 9.50%
 Calcium,min.0.89%max.1.39%
 Phosphorus,min. 0.27%
 Salt,min.0.73%max.1.23%
 Copper,min. max.40ppm
 Selenium,min. 0.62ppm
 Vitamin A,min. 11,000IU/lb

INGREDIENT STATEMENT

Flaked Corn, Soybean Meal, Wheat Middlings, Soybean Hulls, Crimped Barley, Crimped Oats, Cane Molasses, Alfalfa Meal, Calcium Carbonate, Salt, Ammonium Chloride, Monocalcium Phosphate, Brewers Liquid Yeast (*Saccharomyces Cerevisiae*), Potassium Magnesium Sulfate, Ground Limestone, dl-Alpha-Tocopheryl Acetate (Source of Vitamin E), Mineral Oil, Vitamin A Acetate, Cholicalciferol (Source of Vitamin D3), Magnesium Oxide, Copper Sulfate, Selenium Selenite, Lignin Sulfonate, Propionic Acid (a preservative), Acetic Acid, Benzoic Acid (a preservative), Phosphoric Acid, Natural and Artificial Flavoring.

FEEDING DIRECTIONS

Feed as the only source of grain to growing goats, 0.50-1 lb/per/head/day. Feed not more than 0.5lb/hd/day for inactive adult Billies. Supplement with good quality forage, and free choice minerals for all other breeds. Have plenty of clean drinking water available at all times.

CAUTION

Do not allow horses or other equines access to feeds containing Monensin ingestion of Monensin by equines has been fatal. Monensin-medicated feed is intended for use in cattle or in goats. **Consumption by unapproved species may result in toxic reaction.** Do not feed to lactating goats.

MANUFACTURED BY:

Farmers Cooperative Association, Inc.
 820 E South Street Frederick, MD 21701
www.farmerscoop.com
 Net Weight 50 lbs. (22.68Kg.)

3630050

~JU ~PC ~SN

~Bagcount



Medicated 16% Goat Textured