

MoorMan's® IGR Minerals®

To prevent the breeding of horn flies in the manure of treated cattle only. A pesticidally active mineral feed for beef cattle containing methoprene* insect growth regulator for continuous feeding during the fly season.

ACTIVE INGREDIENT:

S-methoprene (CAS Number 65733-16-6)..... 0.02%

OTHER INGREDIENTS: 99.98%
Total 100.00%

GUARANTEED ANALYSIS

Calcium (Ca), not less than.....16.0%
Calcium (Ca), not more than.....19.2%
Phosphorus (P), not less than.....8.0%
Salt (NaCl), not less than13.5%
Salt (NaCl), not more than16.2%
Magnesium (Mg), not less than2.5%
Potassium (K), not less than.....0.2%
Copper (Cu), not less than1100 ppm
Selenium (Se), not less than39 ppm
Zinc (Zn), not less than3800 ppm
Vitamin A, not less than 200,000 International
Units per pound

FEED INGREDIENTS

Monocalcium Phosphate, Dicalcium Phosphate,
Calcium Carbonate, Salt, Defluorinated
Phosphate, Magnesium Oxide, Zinc Oxide,
Manganous Oxide, Copper Sulfate, Ferrous
Sulfate, Mineral Oil, Potassium Sulfate, Potassium
Iodide, Cobalt Carbonate, Vegetable Oil, Cane
Molasses, Plant Protein Products, Iron Oxide,
Calcium Sulfate, Vitamin A Supplement, Vitamin
D3 Supplement, Sodium Selenite, Mineral Oil,
Vitamin E Supplement.

Keep out of reach of children.

CAUTION

Use this product as specified on the label.

EPA REG. NO. 1157-41 EPA Est. 1157-IL-001

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Free-Choice Feeding (self-feeding)

Make sure cattle are not starved for minerals or salt by providing them before feeding MoorMan's IGR Minerals. Then remove salt or mineral supplement and start continuous feeding of MoorMan's IGR Minerals in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity.

The recommended consumption of MoorMan's IGR Minerals is 0.25 lb (1/4 lb) to 0.5 lb (1/2 lb) per 100 lb of body weight per month. Place MoorMan's IGR Minerals near watering or loafing areas and provide one feeder for each 15 to 20 head. Put out only a 5-7 day supply of MoorMan's IGR Minerals at one time and protect it from rain.

If consumption of MoorMan's IGR Minerals is above 0.5 lb (1/2 lb) per 100 lb of body weight per month, reduce the number of feeding stations or relocate feeders in areas frequented less by cattle.

If consumption of MoorMan's IGR Minerals is below 0.25 lb (1/4 lb) per 100 lb of body weight per month, increase the number of feeding stations or relocate in areas more frequented by cattle.

Mixed Ration Feeding (hand-feeding)

When cattle are on supplemental feeds or if mineral consumption is not as recommended, mix MoorMan's IGR Minerals with other non-medicated feeds and feed daily to provide 0.25 lb (1/4 lb) to 0.5 lb (1/2 lb) of MoorMan's IGR Minerals per 100 lb of body weight per month. Start continuous feeding in the spring before horn flies appear on cattle and continue feeding until cold weather restricts horn fly activity.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

18517-QC-TG



7 88399 60001 7

650PCF1K

ADM Animal Nutrition, a division of
Archer Daniels Midland Company, Quincy, IL 62305-3115
Net Weight 50 lb (22.67 kg)